Clinical Development
Biopharma



Immunogenicity assessment of GP2017: A biosimilar to reference adalimumab

Anita Rudy, PhD
Lisbon
February 26, 2019

SANDOZ A Novartis Division


Agenda

1. Introduction
   - Immunogenicity assessment of biosimilars
2. Immunogenicity assessment of GP2017
   - Immunogenicity results from clinical studies
   - Challenges in bioanalytical method validation
3. Summary

Immunogenicity may impact clinical safety and efficacy: assessment needed

- Immunogenicity is a potential concern for **all** biopharmaceuticals
  - Might have an impact on safety and efficacy of the molecule
  - Immunogenicity evaluation is essential

- Many factors can influence immunogenicity, e.g.
  - Route of administration
  - Concomitant medication, disease
  - Aggregates, posttranslational modifications




Immunogenicity assessment of biosimilars: focus on similarity

- Immunogenicity profile of the **molecule per se is already known**
- Focus on confirming that there are **no clinically meaningful differences** between the reference product and the biosimilar
- Head-to-head clinical trials (reference vs. biosimilar) using **state-of-the-art bioanalytical assays**


GP2017 - An approved Sandoz biosimilar to reference adalimumab

- GP2017 is a biosimilar to reference adalimumab
- Up to now GP2017 is approved in US and EU/EEA



- Similar efficacy, safety and immunogenicity demonstrated in confirmatory clinical studies
- Similar PK bioequivalence demonstrated in healthy volunteers
- Comparability demonstrated in comprehensive in vitro studies
- Extensive state of the art characterization demonstrated analytical similarity


Immunogenicity of GP2017 was assessed in a multi-tiered approach



ADA and NAb assay set-up

- State-of-the-art ADA and NAb assays were validated in accordance with current guidelines


ADA assay



GP2017-biotin/sulfo-tag

ADA

NAb assay



GP2017-biotin

NAb

TNF

One vs. two assay approach

1 Assay

Comparability testing during assay validation to show assay detects ADAs against biosimilar and reference equally

- ↑ Conservative approach: detects reliably antibodies against biosimilar
- ↑ No inter-assay variability
- ↑ Blinded study sample analysis possible
- ↑ Wealth of information on reference immunogenicity available
- ↓ ADAs against immunogenic structures (e.g. carbohydrates) unique for reference not detected


2 Assays

Two assays with different properties: different reagents, assay conditions (assay cut-points, sensitivity, stability...)

- ↓ Introduction of additional variability / impact of assay bias on comparative evaluation
- ↓ No blinded study sample analysis possible
- ↓ Data comparison / interpretation more challenging
- ↑ ADAs against potential unique structures of one of the products are detected

One assay approach is suitable for GP2017 immunogenicity assessment


- Specificity demonstrated by immuno-depletion analysis
- Comparable signal inhibition curves



Samples containing fixed ADA concentration were spiked with increasing concentrations of GP2017 or reference product.

PK study to confirm biosimilarity

- Randomized, double-blind, parallel group PK study with three treatment arms to evaluate PK, safety and immunogenicity of GP2017, EU-Reference and US-Reference



ADA development was similar across all three treatment groups


Visit	ADA status	GP2017 N = 107, *n* (%)	EU-Reference N = 106, *n* (%)	US-Reference N = 105, *n* (%)
Day 1[a]	**positive**	2 (1.9)	3 (2.8)	4 (3.8)
Day 16	**positive**	25 (23.4)	27 (25.5)	24 (22.9)
Day 30	**positive**	14 (13.1)	18 (17.0)	16 (15.2)
Day 44	**positive**	19 (17.8)	25 (23.6)	27 (25.7)
Day 72	**positive**	60 (56.1)	70 (66.0)	68 (64.8)

Immunogenicity data in GP2017 PK similarity studies (safety analysis set).
[a] Day 1 samples were collected pre-dose.

von Richter, O. et al., (2019): GP2017, an adalimumab biosimilar: pharmacokinetic similarity to its reference medicine and pharmacokinetics comparison of different administration methods, Expert Opinion on Biological Therapy

Immunogenicity between GP2017 and reference adalimumab is similar

- Immunogenicity is similar, with most ADAs being neutralizing




Overall anti-drug antibody detection rates up to day 72 (safety analysis set).

EU/US-Humira, EU-authorized or US-licensed reference adalimumab; GP2017, Sandoz biosimilar adalimumab.


von Richter, O. et al., (2019): GP2017, an adalimumab biosimilar: pharmacokinetic similarity to its reference medicine and pharmacokinetics comparison of different administration methods, Expert Opinion on Biological Therapy

PK between GP2017 and reference adalimumab is similar

(A) ADA-negative subjects



(B) ADA-positive subjects




Semi-logarithmic serum concentration–time profiles of ADA-negative (A) versus ADA-positive (B) subjects in the PK similarity study (PK analysis set).

Jan Richter, O. et al., (2019): GP2017, an adalimumab biosimilar: pharmacokinetic similarity to its reference medicine and pharmacokinetics comparison of different administration methods, Expert Opinion on Biological Therapy

- Immunogenicity has an impact on PK
- PK in ADA positive and ADA negative subgroups is demonstrated to be similar

Confirmatory safety and efficacy study

- Multicenter, randomized, double-blind, comparator-controlled phase III confirmatory study with four study periods



Blauvelt, A. et al., (2018): Phase III randomized study of the proposed adalimumab biosimilar GP2017 in psoriasis: impact of multiple switches, British Journal of Dermatology

Immunogenicity between GP2017 and reference adalimumab is similar

- Immunogenicity is similar; most ADAs are neutralizing

	Treatment period 1 (randomization to week 17)		Treatment period 2 + extension phase (week 17 to week 51)			
	Reference (*n* = 234)	**GP2017** (*n* = 231)	**Continued Ref** (*n* = 127)	Ref to GP2017 (*n* = 63)	**Continued GP2017** (*n* = 126)	GP2017 to Ref (*n* = 63)
ADA-positive	75/220 **(34.1)**	81/220 **(36.8)**	55/122 **(45.1)**	24/61 (39)	44/123 **(35.8)**	28/60 (47)
NAb-positive	60/75 **(80)**	65/81 **(80)**	47/55 **(85)**	24/24 (100)	38/44 **(86)**	21/28 (75)

Values are the antidrug antibody detection rate, n/N (%), safety analysis set

Modified from: Blauvelt, A. et al., (2018): Phase III randomized study of the proposed adalimumab biosimilar GP2017 in psoriasis: impact of multiple switches, British Journal of Dermatology

Efficacy between GP2017 and reference adalimumab is similar

- PASI 75 response at week 16 in ADA positive and ADA negative subgroups is demonstrated to be similar

ADA status up to week 16	Endpoint	Treatment group	N	n	Adjusted response rate/ SE (%)
Negative	PASI 75 response	GP2017	149	111	74.3 / (3.58)
		Reference	146	105	72.1/ (3.70)
Positive	PASI 75 response	GP2017	40	17	42.8 / (7.72)
		Reference	38	15	39.2 / (7.80)


N- Number of patients per treatment group and subgroup, n- Number of patients per treatment group and subgroup achieving PASI 75 response (per protocol set)

Source: Hyrimoz EPAR


Immunogenicity results cannot be compared to historical data

- Immunogenicity is similar between GP2017 and reference
- Historical data on immunogenicity vary
  - In published studies with adalimumab the proportion of ADA positive subjects varies strongly
  - Differences in patient groups, co-medication
  - Different assay formats
  - No standardization
  - **Assay sensitivity and drug tolerance improved over time**

ADA and NAb assay with high sensitivity and drug tolerance used for immunogenicity assessment of GP2017




ADA assay



NAb assay

Matrix	Validated for different matrices	Validated for different matrices
Assay type	ECL based assay 2-fold acid dissociation	ELISA based CLB assay 2-fold acid dissociation
Sensitivity	~30 ng/mL	~100 ng/mL
Drug tolerance	At least 40µg/mL	At least 40µg/mL
Cut-point	Floating screening Fixed confirmatory	Fixed


Challenges in ADA assay development – screening cut-point



~ 30% of the first ~300 pre-dose samples from clinical study were screened positive

- The assay cut-point i… el of response tha… ble resp… e
  …sing
  …ilable sera
  …exclusion (according to Shankar et al, 2008)
- Floating cut-point factor was determined
- Very low signals


In-study cut-point was determined and used for further analysis



Below 15% of the pre-dose samples from clinical study were screened positive using in-study cut-point

- Highe[illegible]se sample variability
- Slightly higher in-study cut-point determined

Challenges in ADA assay development - LPC





Samples spiked with increasing ADA concentrations were analyzed in the ADA assay.

- LPC: low positive control set to fail in 1% of assay runs
- Calculated LPC below 100 ng/mL was successfully validated

Confirmatory LPC in routine analysis

- LPC above screening cut-point, but LPC could not be confirmed reproducibly
- Confirmatory LPC had to be adapted to a higher concentration





Summary

- Immunogenicity assessment is crucial for **all** biopharmaceuticals
- Similar immunogenicity of GP2017 (approved in EU/EEA and US) to the reference adalimumab was demonstrated
- Sandoz used highly sensitive and highly drug tolerant state-of-the-art assays to determine immunogenicity of GP2017
- Proper cut-point and LPC determination are important for immunogenicity assessment and may be challenging

Acknowledgement

GP2017 Global Team

Clinical Bioanalytics Oberhaching

Thank you

SANDOZ A Novartis Division