

Understanding immune responses to SARS CoV2 infection and vaccination

Alessandro Sette

THE TEAM



Prof. Alex Sette



Alba Grifoni, PhD



Alison Tarke



Ricardo Da Silva Antunes, PhD



Esther Yu Dawan, MD



John Sidney



Prof. Daniela Weiskopf



April Frazier PhD



Prof. Shane Crotty



Camila Coelho, PhD



Zeli Zhang, PhD



Prof. Jen Dan MD/PhD



Jose Mateus, PhD



Nils Methot



Eric Wang



Paul Rubiro

Crotty Lab

Carolyn Rydzynski
Moderbacher
Sydney Ramirez
Prof. Jennifer Dan
Yu 'Alex' Kato
Simon Belanger
Katia Faliti
Christina Kim

Robert Abbott
Eleanor Crotty
Jinyong Choi
Sonya Haupt
Zeli Zhang
Nate Bloom
Ben Goodwin

Sette Lab

Alba Grifoni
Alison Tarke
Esther Yu
Ricardo Antunes
Nils Methot
Jenna Memollo
April Frazier
John Sidney
Paul Rubiro

Weiskopf Lab

Jose Mateus
James Chang

Peters Lab

Bjoern Peters

LJI Bioinformatics Core

Jason Greenbaum

Clinical Studies Core

Gina Levi
Shariza Bautista
Quinn Bui
Jasmine Cardenas

UCSD

Stephen Rawlings
Davey Smith

LJI Flow Cytometry Core

Denise Hinz
Cheryl Kim

JCVI

Richard Scheuermann

Vanderbilt University

Simon Mallal




Philanthropic Support

SARS-CoV-2 should be immunogenic

- Scarce data initially available
- Strong sequence homology and structural similarity with SARS-CoV1
- Similarity also at the epitope level
- Predicted CD4 and CD8 epitopes in SARS CoV2
- Enable experimental testing



Alba Grifoni, PhD




Grifoni et al., Cell Host & Microbe March 2020

Do people develop immunity to SARS CoV2?



Antibodies
(from B cells)

- Important in almost all currently licensed human vaccines



Helper T cells

- Critical for antibody responses
- Protect independent of antibodies in SARS mouse model



Killer T cells

- Important in many viral infections
- Critical to eliminate infected cells


T cell and antibody responses in 'mild' cases of COVID-19

Cell

Article

Targets of T Cell Responses to SARS-CoV-2 Coronavirus in Humans with COVID-19 Disease and Unexposed Individuals

Graphical Abstract



Authors


Alba Grifoni, Daniela Weiskopf, Sydney I. Ramirez, ..., Davey M. Smith, Shane Crotty, Alessandro Sette

Correspondence

shane@lji.org (S.C.), alex@lji.org (A.S.)

In Brief

An analysis of immune cell responses to SARS-CoV-2 from recovered patients identifies the regions of the virus that is targeted and also reveals cross-reactivity with other common circulating coronaviruses




Grifoni et al., Cell May 2020

LJI reagents to measure T cells : shared with 185 labs, 32 different countries, 6 continents and resulted in over 50 collaborative studies

SARS-CoV-2 immune memory following natural infection

- The largest ever study of its kind, for any infection
- Substantial immune memory in most individuals at 8+ months post-COVID
- Immune memory is complex and heterogenous
- T cells, memory B cells, and antibodies have different kinetics
- 5-10% of individuals have low level immune memory at 6+ months



The New York Times

Immunity to the Coronavirus May Last Years, New Data Hint

Blood samples from recovered patients suggest a powerful, long-lasting immune response, researchers reported.




Asst. Prof. Jen Dan
MD/PhD


Dan et al. Science. Jan 2021

Evidence pointing to substantial contributions of T cells to SARS CoV2 immunity


- Early T cell responses correlate with better outcomes and lower viral loads
- CD8 T cells provide control in monkeys
- People lacking antibodies resolve SARS CoV2 infections
- Onset of protection from Moderna or Pfizer vaccines in absence of detectable neutralizing antibodies



Moderbacher et al. Cell Sept 2020




Tan et al, Cell Rep, 2021



Kalimuddin Cell Med, 2021


Vaccine protection against SARS-CoV-2

Protection against
Detectable Infection







Major

Protection against
Hospitalizations & Deaths




Minor



Comparison of immune memory to four COVID-19 vaccines



Characteristic	mRNA-1273 Moderna	BNT162b2 Pfizer	Ad26.COVS.2.S Janssen	NVX-CoV2373 Novavax
Donors, n	30	30	30	12
Gender, n (%)				
Male	12 (40%)	11 (37%)	11 (37%)	6 (60%)
Female	18 (60%)	19 (63%)	19 (63%)	4 (40%)
Age, years (mean ± SD)	43.6 ± 13.9	43.2 ± 17.4	45.6 ± 14.9	38.3 ± 18.0
Race or ethnicity, n				
Asian	3	9	2	0
White	24	16	24	11
Hispanic or latino	9	5	5	1
Days post-vaccination (mean ± SD)				
T2	15 ± 4	15 ± 2	15 ± 3	-
T3	45 ± 6	37 ± 3	47 ± 8	-
T4	105 ± 6	106 ± 4	108 ± 7	120 ± 19
T5	185 ± 9	184 ± 8	185 ± 3	185 ± 11



Comparison of immune memory to four COVID-19 vaccines



Zeli Zhang, PhD





Jen Dan, MD/PhD

PSV, Pseudovirus

Figures modified from Zhang et al. *bioRxiv* 2022.03.18.484953; doi: <https://doi.org/10.1101/2022.03.18.484953>


Comparison of immune memory to four COVID-19 vaccines



Carolyn Rydzynski
Moderbacher, PhD

100% of individuals made memory CD4+ T cells, with Tfh highly represented after mRNA or Novavax vaccination. CD4-CTL were also unexpectedly common

Comparison of immune memory to four COVID-19 vaccines





Jose Mateus, PhD

mRNA vaccines and J&J induced comparable CD8+ T cell frequencies, though memory CD8+ T cells were only detectable in 60-67% of subjects at 6 months

Figures modified from Zhang et al. *bioRxiv* 2022.03.18.484953; doi: <https://doi.org/10.1101/2022.03.18.484953>





Comparison of immune memory to four COVID-19 vaccines



Camila Coelho, PhD


Spike- and RBD-specific memory B cell responses were detected in all individuals to each of the four vaccines, and their frequency increased over time

Comparison of immune memory to four COVID-19 vaccines




What about variants?

SARS-CoV-2



Alpha
(B.1.1.7)



Beta
(B.1.351)



Gamma
(P.1)



B.1.1.519



Kappa
(B.1.617.1)



Delta
(B.1.617.2)



Lambda
(C.37)



R.1



Mu
(B.1.621)



Omicron
(B.1.1.529)


Breadth of SARS CoV2-specific CD4 and CD8 T cell responses

- Identified 280 different CD4 and 523 different CD8 epitopes
- Each individual recognizes multiple epitopes and antigens; conservative estimate of 30-40 epitopes recognized per donor

Tarke et al., Cell Med Rep Jan 2021



- Metanalysis of SARS CoV2 human T cell epitopes; 2049 different human SARS-CoV-2 T cell epitopes defined so far

Grifoni et al., Cell Host & Microbe May 2021



Elemans M, Seich al Basatena N-K, Asquith B (2012) PLoS Comput Biol 8(2): PMID: 22383867

T cell reactivity to SARS-CoV-2 variants in vaccinated individuals




- mRNA-1273
 - BNT162b2
 - Ad26.COVS.2.S
 - NVX-CoV2373
- 2 week post-vax
3-4 month post-vax
6 month post-vax



Alba Grifoni, PhD




Alison Tarke




Tarke A et al, Grifoni and Sette. Cell. Jan 2022 PMID: 35139340

T cell recognition of Omicron is preserved



Tarke A et al, Grifoni and Sette. Cell. Jan 2022 PMID: 35139340



COVID-19 RESPONSE
PRESS BRIEFING

December 28, 2021

CSH Cold Spring Harbor Laboratory
bioRxiv
THE PREPRINT SERVER FOR BIOLOGY

SARS-CoV-2 Vaccination Induces Immunological Memory Able To Cross-Recognize Variants From Alpha to Omicron
A Tarke, A Sette et al.

T cell responses to spike epitopes across SARS-CoV-2 variants, including Omicron, are largely preserved 6 months after vaccination


VACCINES.GOV

- Keeton et al., Nature 2022; [South Africa](#)
- GeurtsvanKessel et al., 2022 Sci Imm; [Netherlands](#)
- Gao et al., Nat Med 2022; [Sweden](#)
- Madelon et al., JAMA Neur 2022; [Switzerland](#)

Tarke A et al, Grifoni and Sette. Cell. Jan 2022 PMID:35139340


Vaccine protection against SARS-CoV-2

Protection against
Detectable Infection




Major

Protection against
Hospitalizations & Deaths






Minor

Waning protection against Detectable Infection




Minimal waning protection against Hospitalizations & Deaths






Unvaccinated




Vaccinated

Recent/
wild type



6 months
Ago/
variants




Boosted



Thoughts on booster vaccination (4th dose) authorized in USA for over 50

Mortality Reduction at Calit Health for Initial Booster and Second Booster




Eric Topol . Ground Truths March 29th

- Israeli studies show positive effects of a 4th immunization in the older age (Pfizer vaccine).
- However:
 - 3 immunizations already protect well from hospitalization and deaths, and short follow up are limitations
 - Antibodies levels restored to peak levels observed after the 3d shot.
 - No measure of T cell and B cell memory.
- Antibody titers contract more rapidly than cellular immunity, which tends to be more stable
- Still need to address long term effects on disease; we might be approaching a stage of limited returns on further immunizations.

Thoughts on booster vaccination (4th dose) authorized in USA for over 50





Mortality Reduction at Calit Health for Initial Booster and Second Booster



Eric Topol . Ground Truths March 29th

- Still there is value in the approval of the 4th immunization
 - new variants can dramatically alter the situation
 - protection offered by the third immunization might wane over time. particularly antibodies in older people.
- The authorization offers an important additional tool to manage the pandemic.
- In the current situation there is a continued need to monitor infections, hospitalizations and variants.

Where are we going? IgG CCC titers are sustained over the years



Ricardo Da Silva
Antunes, PhD




Esther Yu Dawan, MD

Immunological memory to Common Cold Coronaviruses assessed longitudinally over a three-year period. bioRxiv. March 2022. PMID: 35262082

CD4+ T cell responses are also sustained over time


AIM+ (OX40+CD137+)
CD4+ T cells (%)



Days since follow up



Pandemic preparedness. Beyond SARS CoV2

Diagram illustrating NIAID's approach to pandemic preparedness



NIAID's approach to pandemic preparedness. *NIAID*

Reactivity is also detected in non-exposed individuals




- Pre-existing immunity could
 - influence the disease severity of subsequent SARS-CoV-2 infection
 - influence the outcome of SARS-CoV-2 vaccination




Alba Grifoni, PhD

Grifoni et al., Cell May 2020

Widespread evidence of cross-reactive T cell responses within coronaviruses



● SARS-CoV-2	VLKCLKKSLNVAKSE	100
■ 229E	IIKQLKKAMNVAKAE	60
▲ HKU1	QIKQLEKACNIAKSV	47
◆ NL63	LIKQLKRAMNIAKSE	53
◆ OC43	QLKQLEKACNIAKSA	53




● SARS-CoV-2	NVNRFNVAITRAKVG	100
■ 229E	NANRFNVAITRAKKG	87
▲ HKU1	NVNRFNVAITRAKKG	93
▼ MM3-1	NVNRFNLAITRAKKG	87
✱ MM3-3	NVNRFNVAITRARKG	87






Jose Mateus, PhD

Preexisting immune memory effect on low dose Moderna mRNA-1273 COVID-19 vaccine responses



Asst. Prof.
Daniela Weiskopf



Further evidence of cross-reactive T cells roles in prevention of symptomatic COVID-19

RESEARCH ARTICLE | CORONAVIRUS



Cross-reactive CD4⁺ T cells enhance SARS-CoV-2 immune responses upon infection and vaccination

LUCIE LOYAL, JULIAN BRAUN, LARISSA HENZE, BEATE KRUSE, MANUELA DINGELDEY, ULF REIMER, FLORIAN KERN, TATJANA SCHWARZ

MAIKE MANGOLD, CLAUDIA GIESECKE-THIEL, +28 authors, Authors Info & Affiliations

SCIENCE • 8 Oct 2021 • Vol 374, Issue 6564 • DOI: 10.1126/science.abh1823

The Journal of Infectious Diseases

MAJOR ARTICLE



Differential T-Cell Reactivity to Endemic Coronaviruses and SARS-CoV-2 in Community and Health Care Workers

Ricardo da Silva Antunes, Suresh Pallikkuth, Erin Williams, Esther Dawen Yu, Jose Mateus, Lorenzo Quiambao, Eric Wang, Stephen A. Rawlings, Daniel Stadlbauer, Kaijun Jiang, Fatima Amanat, David Arnold, David Andrews, Irma Fuego, Jennifer M. Dan, Alba Grifoni, Daniela Weiskopf, Florian Kramer, Shane Crotty, Michael E. Hoffer, Savita G. Pahwa, and Alessandro Sette



ARTICLE

<https://doi.org/10.1038/s41467-021-27674-x>

OPEN

Cross-reactive memory T cells associate with protection against SARS-CoV-2 infection in COVID-19 contacts

Rhia Kundu, Janakan Sam Narean, Lulu Wang, Joseph Fenn, Timesh Pillay, Nieves Derqui Fernandez, Emily Conibear, Aleksandra Koycheva, Megan Davies, Mica Tolosa-Wright, Seran Hakki, Robert Varro, Eimear McDermott, Sarah Hammett, Jessica Cutajar, Ryan S. Thwaites, Eleanor Parker, Carolina Rosadas, Myra McClure, Richard Tedder, Graham P. Taylor, Jake Dunning, & Ajit Lalvani

Pre-existing polymerase-specific T cells expand in abortive seronegative SARS-CoV-2 infection

Comment on this paper

Leo Swadling, Mariana O. Diniz, Nathalie M. Schmidt, Oliver E. Amin, Aneesh Chandran, Emily Shaw, Corinna Pade, Joseph M. Gibbons, Nina Le Bert, Anthony T. Tan, Anna Jeffery-Smith, Cedric Tan, Christine Y. L. Tham, Stephanie Kucyowicz, Glorienne Aidoo-Micah, Joshua Rosenheim, Jessica Davies, Melanie P. Jensen, George Joy, Laura E McCoy, Ana M Valdes, Lucy van Dorp, Daniel M. Altmann, Rosemary J. Boyton, Charlotte Manisty, Thomas A. Treibel, James C. Moon, COVIDsortium investigators, Francois Balloux, Áine McKnight, Mahdad Noursadeghi, Antonio Bertoletti, Mala K. Maini

JCI The Journal of Clinical Investigation



Recent endemic coronavirus infection is associated with less-severe COVID-19

Manish Sagar, Laura F. White, Joseph P. Mizgerd


J Clin Invest. 2021;131(1):e143380. <https://doi.org/10.1172/JCI143380>.

Concise Communication COVID-19

Graphical abstract



Potential usefulness of cross-reactive HCoV-specific human T cells



- Cross-reactive T cell responses can modulate disease severity
- Several other coronaviruses are also of concern
- Is a "pan-corona" or "pan-sarbeco" vaccine feasible?
- Experimentally determine epitopes/regions that are widely crossreactive
- A T cell vaccine or vaccine component might be effective in broadly preventing severe disease and death



J. Craig Venter™
I N S T I T U T E




Alba Grifoni, PhD



Alison Tarke

Thoughts on broadly crossreactive T cells

- T cell inducing vaccine components as a broad concept to enhance preparedness against future possible pandemics
- Strategy could be considered for virus families of pandemic preparedness concern (Arenav, Flavi, Bunya, Paramixo, Togav, Picorna and Filioviridae)
- Not an alternative to antibody strategies, but rather synergistic
- The human influenza T cell repertoire is broad and multispecific
- Human epitopes immunogenic conserved in Old and New world Arenaviruses
- Cross-reactivity of epitopes from multiple DENV serotypes and ZIKV




NIAID's approach to pandemic preparedness. NIAID


Overall Conclusions

- SARS CoV2 infection induces a multispecific and multifunctional adaptive responses which is durable over the 6-8 month period
- Neutralizing antibodies likely key to protect from infection, but substantial contributions of T cells likely at the level of protection from severe disease
- Side-by-side longitudinal comparison of different vaccine platforms reveals durable B and T cell responses, and allow correlation with vaccine efficacy
- T cell reactivity is largely preserved at the level of SARS CoV2 variants, including Omicron and Delta
- CCC cross-reactivity albeit rare, is detected and has biological relevance, with potential pandemic preparedness implications
- Responses to CCC are sustained in the adult population, offering a potential glimpse in endemicity


Protection against detectable infection





Protection against hospitalizations and deaths



Waning protection against detectable infection



Minimal waning protection against hospitalizations and deaths



Adapted from Kaiser Study. (Tartof et al. Lancet 2021. PMID 34619098)